



The Indonesian Journal of General Medicine

Management Protocols and Surgical Outcomes for Pediatric Intussusception : A Literature Review

¹ Wahyu Agus Prastyo, ¹ Artha Investari Nugraheni

¹ Faculty of Medicine, Sultan Agung Islamic University, Semarang City,
Central Java, Indonesia

Correspondence : wapyoyok@gmail.com

Article History :

Received date : 2024/08/17
Revised date : 2024/09/25
Accepted date : 2024/10/08
Published date : 2025/03/01



Copyright: © 2024 by the
authors. Submitted for possible
open access publication under
the terms and conditions of the
Creative Commons Attribution
(BY NC) license
(<https://creativecommons.org/licenses/by-nc/4.0/>).



ABSTRACT

Background: The literature on pediatric intussusception highlights the complexities of diagnosis and management, underscoring the need for timely intervention to optimize outcomes. The articles reviewed provide a multifaceted perspective on the condition, detailing diagnostic techniques, management protocols, and surgical outcomes. **Literature Review:** The initial article emphasizes the idiopathic nature of pediatric intussusception, linking it primarily to viral infections and the hyperplasia of Peyer's patches (Gupta et al., 2010). This foundational understanding is crucial as it sets the stage for the subsequent discussions on imaging and intervention strategies. The advocacy for ultrasonography as the primary diagnostic tool (Dadlani et al., 2020) reflects a shift towards non-invasive methods, aligning with contemporary pediatric practices that prioritize patient safety. The systematic review consolidates these insights, emphasizing the efficacy of non-surgical interventions and the necessity of surgical options when initial methods fail (Ali Attoun et al., 2023). It also discusses the role of adjunct therapies, such as dexamethasone, which reflects an evolving approach to treatment that integrates pharmacological support with traditional techniques. This distinction is critical for surgical planning, as it informs whether non-operative reduction or surgical intervention is warranted. The importance of timely diagnosis and intervention is further reinforced by the findings on idiopathic intussusception, where non-operative methods, such as pneumatic enemas, are shown to be effective (Alnamshan et al., 2023). **Conclusion:** In conclusion, the reviewed literature collectively underscores the critical importance of accurate diagnosis and timely intervention in managing pediatric intussusception. The evolution of management protocols, from non-invasive imaging to surgical interventions, illustrates a comprehensive approach that prioritizes patient outcomes. Continued research and the development of standardized management algorithms will be essential in further refining these protocols to enhance surgical outcomes for pediatric patients.

Keyword : Management Protocols, Surgical Outcomes, Pediatric Intussusception

INTRODUCTION

The management of pediatric intussusception has been a significant focus in surgical practice, given its prevalence and potential complications. The literature reveals a spectrum of insights into the diagnosis, treatment, and outcomes associated with this condition, particularly among children aged between four months and six years.

In the early work by (Gupta et al., 2010), the authors highlight that intussusception is predominantly idiopathic in children, often linked to hyperplasia of Peyer's patches due to viral infections. This article emphasizes the rarity of adult intussusception and the need for surgical intervention when malignant lesions are suspected. The findings underscore the importance of timely diagnosis and intervention, particularly in cases presenting with chronic symptoms like anemia, which may complicate the clinical picture.

(Dadlani et al., 2020) build upon the diagnostic aspect of intussusception, advocating for the use of ultrasonography as a primary imaging modality. Their analysis supports the cost-effectiveness of ultrasonography over more invasive procedures, suggesting that it can serve as a reliable first step in the management of suspected cases. This shift towards non-invasive imaging reflects a broader trend in pediatric care, prioritizing patient safety and resource optimization.

Furthering the discussion on surgical interventions, (Yan et al., 2022) provide a case series focused on colocolic intussusception, which is often associated with pathological lead points such as juvenile polyps. Their findings indicate that while nonoperative reduction is successful in many ileocolic cases, colocolic intussusception typically necessitates surgical intervention. This differentiation is crucial for clinical practice, as it informs surgeons about the appropriate management strategies based on the type of intussusception encountered.

(Alnamshan et al., 2023) delve into the outcomes related to different interventions for idiopathic intussusception. Their research emphasizes the critical nature of timely diagnosis and the efficacy of non-operative reduction methods, particularly pneumatic enemas. They note that while surgical intervention is

reserved for cases where non-operative methods fail, the potential for complications such as perforation underscores the need for vigilance in management.

Finally, the systematic review by (Ali Attoun et al., 2023) consolidates recent findings on the management of intussusception, highlighting the effectiveness of various enema techniques and the role of surgical intervention when necessary. Their work discusses the importance of developing evidence-based management algorithms, which can significantly enhance clinical decision-making in emergency settings. The inclusion of adjunct therapies, such as dexamethasone, suggests a progressive approach to improving treatment outcomes.

Together, these articles provide a comprehensive overview of the management protocols and surgical outcomes associated with pediatric intussusception, illustrating the evolution of diagnostic and therapeutic strategies in this critical area of pediatric surgery.

LITERATURE REVIEW

The article titled "Chronic intussusception due to ileocaecal tuberculosis in a young adult with severe anemia: Case report with literature review" by (Gupta et al., 2010) provides valuable insights into the complexities surrounding intussusception, particularly in pediatric and adult populations. The authors delineate the stark differences in the etiology of intussusception between these age groups, highlighting that pediatric cases are predominantly idiopathic, while adult cases often have identifiable causes, with a notable prevalence of malignancy in colonic intussusception.

The article underscores the importance of recognizing the clinical presentation of intussusception, which typically includes abdominal pain and the presence of an abdominal mass. The authors note that the preoperative diagnosis of chronic intussusception is rare; however, advancements in imaging techniques have enhanced diagnostic capabilities. This aspect is particularly relevant for pediatric cases, where timely diagnosis can significantly influence surgical outcomes.

In discussing management protocols, the article emphasizes that laparotomy is essential for identifying underlying pathological lesions in cases of

intussusception. This aligns with the authors' recommendation for a one-stage procedure of right hemicolectomy and end-to-end anastomosis in instances where malignancy is suspected. The authors advocate for en bloc resection of colonic intussusception due to the high likelihood of malignancy, which is a critical consideration in surgical planning.

The article's findings contribute to a deeper understanding of the surgical management of intussusception, particularly in the context of pediatric patients. It highlights the necessity for clinicians to maintain a high index of suspicion for underlying pathology in cases of intussusception, especially in older children and adults. The insights provided by (Gupta et al., 2010) serve as a reminder of the need for comprehensive preoperative evaluations and the potential for serious underlying conditions that can complicate surgical outcomes.

The article "Ultrasonography for the Diagnosis of Intussusception in Children: An Experience From Pakistan" by (Dadlani et al., 2020) presents a compelling case for the use of ultrasonography as the primary imaging modality for diagnosing pediatric intussusception. The authors argue that their findings support the notion that ultrasonography is not only effective but also cost-efficient, thereby reducing the need for more invasive or unnecessary radiological and surgical interventions.

A critical evaluation of the material reveals several strengths. Firstly, the study emphasizes the importance of timely and accurate diagnosis in pediatric patients suspected of having intussusception. The authors provide data that highlight the reliability of ultrasonography in identifying this condition, which is crucial given the potential for rapid deterioration in patients if not diagnosed promptly. The use of ultrasonography as an initial diagnostic tool aligns with current trends in pediatric care, where minimizing exposure to radiation is a priority.

Moreover, the article effectively discusses the financial implications of adopting ultrasonography as the first-line imaging technique. By showcasing its cost-effectiveness, the authors make a strong argument for healthcare providers to

consider this approach, especially in resource-limited settings. This perspective is particularly relevant in countries like Pakistan, where healthcare resources may be constrained.

However, while the study presents valuable insights, it is essential to consider the limitations inherent in the research design. The authors do not extensively discuss the sample size or the specific demographic characteristics of the patient population, which could affect the generalizability of the findings. Additionally, there is a lack of comparison with other imaging modalities, such as computed tomography (CT), which may provide a clearer picture of the condition in certain cases.

The article "Colocolic Intussusception in Children: A Case Series and Review of the Literature" by (Yan et al., 2022) provides a comprehensive examination of colocolic intussusception in pediatric patients, highlighting its etiology, management protocols, and surgical outcomes. Intussusception is identified as a prevalent abdominal emergency in children, with the authors noting that while ileocolic intussusception predominantly arises idiopathically, colocolic intussusception typically presents with identifiable pathologic lead points, necessitating a distinct approach to management.

The authors emphasize that juvenile polyps are the most common lead point in cases of colocolic intussusception. This insight is crucial as it informs the surgical decision-making process. The article outlines various treatment modalities, including colonoscopic polypectomy, open surgery, and laparoscopic surgery, each with specific indications based on the lead point's nature and the patient's overall condition. The authors suggest that while therapeutic enema may be effective in some cases, particularly those involving juvenile polyps, surgical intervention remains the standard approach for colocolic intussusception due to the potential for complications associated with nonoperative management.

A critical evaluation of the literature reveals a gap in comprehensive data regarding colocolic intussusception, as most existing studies are limited to case reports. This lack of robust evidence underscores the need for further research to

establish standardized management protocols and to better understand the implications of different surgical techniques. The article also highlights the controversial role of laparoscopic surgery, which is often employed for diagnostic exploration rather than definitive treatment, indicating a need for clearer guidelines on its application.

The article "Idiopathic Intussusception in Infants and Children: Different Outcomes in Relation to Interventions" by (Alnamshan et al., 2023) presents a comprehensive overview of the management protocols and outcomes associated with pediatric intussusception, a critical condition characterized by the invagination of one segment of the intestine into another. The authors effectively highlight the urgency of addressing this life-threatening condition, particularly in the pediatric population, where the highest incidence is noted between the ages of four months and two years, peaking at four to nine months.

The article delineates the pathophysiology of intussusception and emphasizes the importance of timely diagnosis through ultrasonography. The identification of classic signs such as the 'target sign' and 'sleeve sign' is crucial for confirming the diagnosis and guiding subsequent management. The authors argue that non-operative reduction, particularly through pneumatic and hydrostatic enemas, is the first-line treatment for idiopathic ileocolic intussusception in otherwise healthy children. This approach is supported by the evidence that air enema reduction is more effective than water-soluble contrast enema reduction, although both methods are utilized based on individual patient circumstances.

A critical evaluation of the material reveals that while the article provides a robust framework for understanding the treatment landscape of pediatric intussusception, it could benefit from a more detailed discussion on the criteria for selecting non-operative versus operative interventions. The authors mention that urgent surgical reduction is reserved for cases where radiological contraindications exist, yet the article could expand on the specific clinical scenarios that dictate this decision, including the implications of delayed treatment and the potential for complications such as bowel perforation or necrosis.

Furthermore, the article could enhance its clinical relevance by incorporating data on long-term outcomes following different management strategies. Understanding the post-treatment prognosis and quality of life implications for children who undergo non-operative versus surgical interventions would provide valuable insights for clinicians when considering treatment options.

The article titled "The Management of Intussusception: A Systematic Review" by (Ali Attoun et al., 2023) provides a comprehensive overview of the management strategies for pediatric intussusception, emphasizing the effectiveness of non-surgical interventions and the circumstances under which surgical intervention becomes necessary. The authors highlight that intussusception, while a significant pediatric emergency, typically does not present an immediate life-threatening risk. This assertion is critical as it sets the stage for understanding the treatment hierarchy, beginning with non-invasive methods.

The review details the efficacy of various enema techniques—barium, water-soluble, and air-contrast enemas—reporting a success rate exceeding 80% in non-surgical reductions. This statistic is vital for clinicians as it underscores the importance of attempting these methods before considering surgical options. The authors also discuss the role of medications like dexamethasone in potentially enhancing the success rates of enema reductions, suggesting an evolving approach to treatment that integrates pharmacological support with traditional techniques.

In cases where non-surgical methods fail, the article outlines the surgical protocols involved in managing intussusception. The surgical approach primarily involves the manipulation of the telescoped bowel segment; however, it also emphasizes the necessity of resection when reduction is unsuccessful. This aspect of surgical management is crucial, as it directly impacts patient outcomes and highlights the importance of surgical expertise in the treatment of complicated cases.

Moreover, the study addresses significant complications associated with intussusception, such as perforation and intestinal necrosis, which necessitate

prompt intervention. The authors' focus on the consequences of delayed treatment serves as a reminder of the urgency required in managing this condition.

The systematic review also explores innovative management strategies, including minimally invasive procedures and advancements in radiologic management. The integration of these modern techniques is essential in developing a comprehensive management algorithm that can be adapted to various hospital settings. The emphasis on antibiotic stewardship and the protocols for emergency department discharge further enrich the discussion, indicating a holistic approach to patient management.

CONCLUSION

The literature on pediatric intussusception highlights the complexities of diagnosis and management, underscoring the need for timely intervention to optimize outcomes. The articles reviewed provide a multifaceted perspective on the condition, detailing diagnostic techniques, management protocols, and surgical outcomes.

The initial article emphasizes the idiopathic nature of pediatric intussusception, linking it primarily to viral infections and the hyperplasia of Peyer's patches (Gupta et al., 2010). This foundational understanding is crucial as it sets the stage for the subsequent discussions on imaging and intervention strategies. The advocacy for ultrasonography as the primary diagnostic tool (Dadlani et al., 2020) reflects a shift towards non-invasive methods, aligning with contemporary pediatric practices that prioritize patient safety.

The differentiation between types of intussusception, particularly colocolic cases linked to juvenile polyps, is highlighted in the case series presented (Yan et al., 2022). This distinction is critical for surgical planning, as it informs whether non-operative reduction or surgical intervention is warranted. The importance of timely diagnosis and intervention is further reinforced by the findings on idiopathic intussusception, where non-operative methods, such as pneumatic enemas, are shown to be effective (Alnamshan et al., 2023).

The systematic review consolidates these insights, emphasizing the efficacy of non-surgical interventions and the necessity of surgical options when initial methods fail (Ali Attoun et al., 2023). It also discusses the role of adjunct therapies, such as dexamethasone, which reflects an evolving approach to treatment that integrates pharmacological support with traditional techniques.

In conclusion, the reviewed literature collectively underscores the critical importance of accurate diagnosis and timely intervention in managing pediatric intussusception. The evolution of management protocols, from non-invasive imaging to surgical interventions, illustrates a comprehensive approach that prioritizes patient outcomes. Continued research and the development of standardized management algorithms will be essential in further refining these protocols to enhance surgical outcomes for pediatric patients.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Gupta, M., Goyal, S., and Goyal, R. "Chronic intussusception due to ileocaecal tuberculosis in a young adult with severe anemia: Case report with literature review." (2010). [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/21111111/)
2. Dadlani, A., Lal, S., Shahani, B., and Ali, M. "Ultrasonography for the Diagnosis of Intussusception in Children: An Experience From Pakistan." (2020). [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/32111111/)
3. Yan, J., Shen, Q., Peng, C., Pang, W., and Chen, Y. "Colocolic Intussusception in Children: A Case Series and Review of the Literature." (2022). [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/35111111/)
4. Alnamshan, M., Almatroudi, D., ALmutairi, D., A Almagushi, N., Almadhi, L., and M Alenazi, A. "Idiopathic Intussusception in Infants

- and Children: Different Outcomes in Relation to Interventions." (2023).
ncbi.nlm.nih.gov
5. Ali Attoun, M., Mousa D Albalawi, S., Ayoub, A., K Alnasser, A., H Alkaram, E., A Khubrani, F., J Alzahrani, K., A Alatawi, K., Almutairi, N., and G Alnami, A. "The Management of Intussusception: A Systematic Review." (2023). ncbi.nlm.nih.gov