



The Indonesian Journal of General Medicine

Prevalence of Depression in Patients With Mild Cognitive Impairment

¹ Muhammad Iqbal, ² Muhammad Imam

¹ Faculty of Medicine, General Achmad Yani University, Cimahi City, West
Java, Indonesia

² Faculty of Medicine, Padjadjaran University, Bandung City, West Java,
Indonesia


Correspondence : dr.iqbalmuhammad2891@gmail.com

Article History :

Received date : 2024/04/12
Revised date : 2024/05/22
Accepted date : 2024/06/19
Published date : 2024/07/02



Copyright: © 2024 by the
authors. Submitted for possible
open access publication under
the terms and conditions of the
Creative Commons Attribution
(BY NC) license
(<https://creativecommons.org/licenses/by-nc/4.0/>).



ABSTRACT

Background: The literature on the prevalence of depression in patients with mild cognitive impairment (MCI) reveals a complex interplay between these conditions that significantly affects patient outcomes. Various studies illustrate how depressive symptoms can exacerbate cognitive decline, increase the risk of progression to dementia, and complicate diagnosis and treatment strategies. **Literature Review:** The examination of depressive symptoms in patients with MCI, particularly in the context of Parkinson's disease, indicates that these symptoms significantly impair recognition performance. The relationship between social engagement and depression is also critical, as increased social support may mitigate depressive symptoms and improve overall well-being. The challenges in differentiating between MCI and depression are underscored by the overlapping cognitive impairments, suggesting that subsyndromal depressive symptoms may serve as early indicators of dementia. The bidirectional association between depression and cognitive impairment is particularly pronounced in primary care settings, where depression rates among individuals with MCI can reach as high as 32%. **Conclusion:** In conclusion, the literature collectively highlights the urgent need for integrated approaches in the assessment and management of depression and MCI. Recognizing the interplay between these conditions can lead to improved outcomes through early detection, targeted interventions, and a holistic approach to patient care.

Keyword: Prevalence, Depression, Mild Cognitive Impairment

INTRODUCTION

The prevalence of depression in patients with mild cognitive impairment (MCI) has garnered significant attention in recent years, highlighting the complex interplay between these two conditions. The literature reveals a troubling association, with depressive symptoms often exacerbating cognitive decline and increasing the risk of progression to more severe forms of dementia. In a study by (Lara Soares Blum Malak et al., 2017), the authors found that patients with mild cognitive impairment associated with Parkinson's disease exhibited impaired recognition performance, suggesting that depressive symptoms play a critical role in cognitive decline. The findings indicated that recognition memory impairment is a predictive factor for dementia, emphasizing the need for early detection and intervention in this vulnerable population.

(Said, 2018) further explored the mental health difficulties accompanying subjective cognitive complaints and MCI, revealing a significant association between depression, anxiety, and cognitive

decline. The review underscored the necessity for standardized measurement of subjective cognitive complaints (SCC) and the effects of mental health treatment on cognitive outcomes, indicating that common mental health difficulties could substantially impact the trajectory of cognitive decline.

In a community-dwelling context, (Song et al., 2019) identified low-level physical activity as a significant contributor to depressive symptoms in older adults with MCI. The study highlighted the importance of maintaining social engagement and physical activity to mitigate depressive symptoms, thereby improving overall well-being and potentially slowing cognitive decline. The relationship between social isolation and depression was particularly emphasized, suggesting that interventions aimed at increasing social support could be beneficial.

(Han et al., 2021) conducted a longitudinal study that established a dose-response relationship between depressive symptoms and the incidence of MCI, particularly among older adults and women. Their

findings reinforce the idea that addressing depressive symptoms in these populations may be crucial in preventing the onset of MCI.

(Cankaya et al., 2024) examined the challenges in differentiating between MCI and depression, noting the overlapping cognitive impairments and the role of hippocampal integrity. Their work suggests that subsyndromal depressive symptoms might serve as early indicators of dementia, complicating diagnosis and treatment strategies. This underscores the necessity for comprehensive assessments to accurately identify patients at risk.

Additionally, (Xue et al., 2024) investigated biomarkers related to synaptic function and cognitive impairment in depression, further supporting the notion that cognitive deficits are prevalent among individuals with depressive disorders. Their findings contribute to the understanding of how depression may predict the occurrence of MCI.

Finally, (D. Federman et al., 2024) highlighted the bidirectional association between depression and

cognitive impairment in a primary care setting, revealing that depression rates among individuals with MCI can reach as high as 32%. The study emphasizes the clinical relevance of recognizing these conditions in tandem, as they share common pathophysiological pathways and significantly affect patient care.

Collectively, these studies illuminate the intricate relationship between depression and mild cognitive impairment, highlighting the urgent need for integrated approaches in assessment and management to improve outcomes for affected individuals.

LITERATURE REVIEW

The article "Symptoms of depression in patients with mild cognitive impairment in Parkinson's disease" by (Lara Soares Blum Malak et al., 2017) provides a comprehensive examination of the interplay between depression and mild cognitive impairment (MCI) within the context of Parkinson's disease (PD). The authors highlight that MCI cannot be solely defined by deficits in frontal executive function, as previous literature might suggest.

Instead, they propose that recognition performance is significantly impacted by the presence of depressive symptoms, indicating a complex relationship between cognitive decline and mood disorders.

One of the critical insights from this study is the observation that patients with both MCI and depression (PDDS group) exhibited impaired recognition performance compared to those without depressive symptoms (CGDS group). This finding emphasizes the detrimental effect of depression on cognitive resources, which are already compromised in individuals with MCI. The authors argue that cognitive decline in depressed patients is particularly pronounced in episodic memory, a crucial domain for daily functioning and quality of life.

Moreover, the article underscores the importance of recognizing the combined effects of depression and MCI, as this dual diagnosis significantly heightens the risk of conversion to dementia. The authors present evidence that impairment in recognition memory serves as a predictive factor for

dementia, reinforcing the need for early detection and intervention strategies. This is particularly relevant given that both conditions adversely affect autonomy and independence, leading to increased rates of institutionalization and mortality.

The study also identifies common depressive symptoms among the PD group, including difficulties with work, fatigue, and sleep disorders. These symptoms not only contribute to the cognitive decline observed but also exacerbate the overall quality of life for these patients. By highlighting the specific cognitive domains most affected—namely episodic memory and attention—the authors provide a clear framework for understanding how depression complicates the clinical picture of MCI in Parkinson's disease.

The article "Exploring fear of dementia, subjective cognitive complaints and common mental health difficulties in screening for Mild Cognitive Impairment" by (Said, 2018) provides a comprehensive examination of the interplay between mental health issues, particularly depression and

anxiety, and the progression of cognitive decline in individuals with Mild Cognitive Impairment (MCI). The review synthesizes findings from 17 studies, highlighting a notable association between common mental health difficulties and cognitive decline, as identified in 81% of the included papers.

Said's review critically evaluates the relationship between subjective cognitive complaints (SCC) and mental health challenges, underscoring the prevalence of depression among individuals presenting with MCI. The article suggests that mental health difficulties are not merely co-occurring conditions but may significantly influence the trajectory of cognitive decline. This perspective is crucial for clinicians, as it emphasizes the need for a holistic approach to assessing and treating patients with MCI, where mental health is considered an integral component of cognitive health.

One of the notable strengths of the article is its identification of significant variability in the measurement of SCC across the

studies reviewed. This variability raises important considerations regarding the interpretation of findings within the literature and the implications for clinical practice. The inconsistency in measurement tools may lead to divergent conclusions about the prevalence and impact of depression and anxiety in patients with MCI. Therefore, Said advocates for the standardization of SCC measurement, which could enhance the reliability of research outcomes and improve clinical assessments.

Furthermore, the article calls for future research to evaluate the effects of interventions targeting depression and anxiety on cognitive decline. This recommendation is particularly pertinent, as it opens avenues for exploring therapeutic strategies that could mitigate the cognitive deterioration associated with MCI. By addressing mental health difficulties, there is potential not only to improve the quality of life for individuals with MCI but also to slow the progression towards dementia.

The article "Identifying the Factors Related to Depressive

Symptoms Amongst Community-Dwelling Older Adults with Mild Cognitive Impairment" by (Song et al., 2019) provides a comprehensive examination of the interplay between mild cognitive impairment (MCI) and depressive symptoms in older adults. The authors emphasize the significant role that low-level physical activity plays in the prevalence of depressive symptoms among this demographic, aligning with a growing body of literature that highlights the beneficial effects of physical activity on mood and mental health.

The article effectively articulates how physical activity can stimulate the release of neurotransmitters such as serotonin and dopamine, which are vital for enhancing mood and promoting a sense of well-being. This biochemical perspective is crucial as it not only underscores the physiological benefits of exercise but also suggests potential therapeutic avenues for managing depression in individuals with MCI. Furthermore, the authors discuss the social withdrawal hypothesis, positing that increased sedentary behavior may lead to reduced social interactions, thereby

elevating the risk of depression. This insight is particularly relevant as it connects physical activity levels with social engagement, illustrating how lifestyle choices can significantly impact mental health outcomes.

A critical evaluation of the article reveals its thorough approach to identifying at-risk individuals. The authors advocate for the incorporation of subjective assessments of disease-related factors in identifying those with MCI who may be vulnerable to depressive symptoms. This perspective is vital, as it underscores the necessity of a holistic approach to treatment that considers both psychological and social dimensions of health. The strong correlation presented between social networks, social support, and depressive symptoms further emphasizes the importance of fostering social connections as a protective factor against depression.

However, while the article presents compelling evidence regarding the relationship between physical activity, social engagement, and depressive symptoms, it could benefit from a more detailed

exploration of specific interventions that could be implemented to enhance physical activity and social interaction among older adults with MCI. Additionally, the article could have expanded on the mechanisms by which social support influences mood and the potential barriers to engaging in physical activity within this population.

The article titled "Depressive symptoms and cognitive impairment: A 10-year follow-up study from the Survey of Health, Ageing and Retirement in Europe" by (Han et al., 2021) presents a compelling examination of the relationship between depressive symptoms and the incidence of mild cognitive impairment (MCI). The authors utilize a longitudinal approach to analyze data collected over a decade, offering valuable insights into how depressive symptoms may contribute to cognitive decline, particularly in older adults.

One of the central findings of the study is the identification of a significant association between depressive symptoms and the incidence of MCI, characterized by a

dose-response relationship. This indicates that as the severity of depressive symptoms increases, so does the likelihood of developing MCI. This finding is particularly pronounced among individuals aged 70 years and older, as well as among women, suggesting that these demographics may be at heightened risk. The authors argue that this relationship necessitates targeted interventions aimed at treating depressive symptoms within these groups, which could potentially serve as a preventive measure against the onset of MCI.

The methodology employed in the study is robust, utilizing a large sample size and a comprehensive follow-up period that enhances the reliability of the findings. The use of standardized measures for both depressive symptoms and cognitive function allows for a nuanced understanding of the interplay between these two constructs. However, the article could benefit from a more in-depth discussion regarding the potential mechanisms underlying the observed associations. For instance, exploring how neurobiological changes associated

with depression might influence cognitive processes would enrich the analysis.

Furthermore, while the authors highlight the importance of addressing depressive symptoms to mitigate the risk of MCI, they do not extensively discuss the implications of such interventions. A more detailed exploration of specific treatment modalities and their effectiveness in this context would provide practical guidance for clinicians and caregivers.

The article "Differentiating MCI from depression through verbal memory scores" by (Cankaya et al., 2024) provides a nuanced examination of the intricate relationship between Mild Cognitive Impairment (MCI) and depression, particularly emphasizing the role of hippocampal integrity in memory performance. The authors present a compelling argument that structural and functional changes in the hippocampus are pivotal in understanding the interplay between these two conditions. The evidence presented suggests that depression not only serves as a risk factor for

dementia but also complicates the diagnostic landscape of MCI due to overlapping symptoms and cognitive deficits.

The article highlights that the prevalence of depression among individuals with MCI is notably significant, with meta-analyses indicating a rate of 32%. This statistic underscores the importance of recognizing depressive symptoms as potential early indicators of dementia, a critical insight for both clinical practice and research. The authors argue convincingly that subsyndromal depressive symptoms can manifest during the prodromal stages of dementia, complicating the diagnostic process. This assertion is particularly relevant given the challenges in distinguishing between MCI and major depressive disorder (MDD), as cognitive impairments in MCI may not be as pronounced as those in full-blown dementia.

Furthermore, the article discusses the anatomical and functional overlaps in brain regions responsible for both cognitive and emotional processing. This overlap provides a robust rationale for the

increased dementia risk in patients with MDD who also exhibit cognitive impairment. The authors emphasize the necessity for regular cognitive assessments in these patients to facilitate early identification and intervention, which is crucial for improving patient outcomes.

Despite these valuable insights, the article also acknowledges the challenges inherent in accurately diagnosing MCI in the presence of depressive symptoms. The authors call for a more comprehensive and specific assessment approach, which is essential for distinguishing between MCI and MDD. The Turkish Verbal Memory Process Test (VMPT) is highlighted as a useful tool for this purpose, given its focus on eliciting verbal information and assessing executive functions.

The article titled "Synapse-Related Serum and P300 Biomarkers Predict the Occurrence of Mild Cognitive Impairment in Depression" by (Xue et al., 2024) presents a significant exploration into the intersection of depression and mild cognitive impairment (MCI). The

authors aim to elucidate the relationship between these two conditions, emphasizing that cognitive impairment frequently coexists with depression. This study is particularly relevant in the context of understanding the prevalence of depression in patients with MCI.

The authors methodically investigate the predictive capabilities of synapse-related serum biomarkers and P300 event-related potentials in identifying individuals who may develop MCI as a consequence of depression. Their findings suggest a notable correlation between the biomarkers and the cognitive deficits observed in depressed patients, thereby reinforcing the notion that depression is not merely a psychological condition but can manifest with substantial cognitive repercussions.

A critical evaluation of the material reveals that the study is grounded in a robust methodology, utilizing both biochemical and neurophysiological measures to assess cognitive function in the context of depression. By employing serum biomarkers, the authors

provide an innovative approach to identifying at-risk populations for MCI, which could have profound implications for early intervention strategies. Furthermore, the use of P300 as a measure of cognitive processing adds depth to the analysis, as it reflects the underlying neural mechanisms associated with cognitive impairment.

However, while the study presents compelling evidence linking depression with MCI, it is important to consider the limitations inherent in the research design. The sample size and demographic diversity may affect the generalizability of the findings. Additionally, the cross-sectional nature of the study limits the ability to draw causal inferences between depression and the development of MCI. Longitudinal studies would be beneficial to further substantiate the predictive relationship highlighted by the authors.

The article titled "Relationship Between Cognitive Impairment and Depression Among Middle Aged and Older Adults in Primary Care" by (D. Federman et al., 2024) provides a comprehensive

examination of the interrelationship between cognitive impairment and depression, particularly in a primary care setting. The authors highlight a significant prevalence of depression among adults, ranging from 8% to 20%, while noting an alarming rate of 32% among individuals with mild cognitive impairment (MCI) and even higher rates (37% to 41%) in those with dementia. This underscores the critical need for clinicians to be vigilant in recognizing depression in patients exhibiting cognitive difficulties.

One of the key insights from the study is the bidirectional association between depression and cognitive impairment. The authors argue that depression not only affects cognitive performance—manifesting as difficulties with concentration and decision-making—but also poses a risk for cognitive decline and accelerates the progression from MCI to dementia. This relationship is particularly concerning for older adults, as the presence of depression can complicate the management of cognitive impairment and exacerbate the overall health burden.

The authors conducted their research with a focus on identifying previously undiagnosed cognitive impairment in older adults presenting with comorbid depression in primary care settings. This approach is clinically relevant, as it emphasizes the importance of screening for cognitive impairment in patients diagnosed with depression, which may often go overlooked. The implications of these findings are profound, suggesting that early identification and treatment of depression in older adults with MCI could potentially mitigate cognitive decline and improve quality of life.

CONCLUSION

The literature on the prevalence of depression in patients with mild cognitive impairment (MCI) reveals a complex interplay between these conditions that significantly affects patient outcomes. Various studies illustrate how depressive symptoms can exacerbate cognitive decline, increase the risk of progression to dementia, and complicate diagnosis and treatment strategies.

The examination of depressive symptoms in patients with MCI, particularly in the context of Parkinson's disease, indicates that these symptoms significantly impair recognition performance ((Lara Soares Blum Malak et al., 2017)). This impairment not only affects cognitive function but also serves as a predictive factor for dementia, highlighting the critical need for early detection and intervention. Furthermore, the interplay between subjective cognitive complaints and mental health difficulties, particularly anxiety and depression, emphasizes the necessity for standardized measures to assess these variables ((Said, 2018)). The findings suggest that mental health interventions could substantially impact cognitive outcomes for individuals with MCI.

In community-dwelling older adults, low levels of physical activity have been identified as a significant contributor to depressive symptoms, with implications for cognitive health ((Song et al., 2019)). The relationship between social engagement and depression is also critical, as increased social support may mitigate depressive symptoms and improve

overall well-being. Additionally, a longitudinal study established a dose-response relationship between depressive symptoms and the incidence of MCI, particularly among older adults, indicating the importance of addressing depression as a preventive measure ((Han et al., 2021)).

The challenges in differentiating between MCI and depression are underscored by the overlapping cognitive impairments, suggesting that subsyndromal depressive symptoms may serve as early indicators of dementia ((Cankaya et al., 2024)). This complexity necessitates comprehensive assessments to accurately identify at-risk patients. Moreover, research into synaptic function and biomarkers related to cognitive impairment in depression further supports the notion that cognitive deficits are prevalent among individuals with depressive disorders ((Xue et al., 2024)).

The bidirectional association between depression and cognitive impairment is particularly pronounced in primary care settings,

where depression rates among individuals with MCI can reach as high as 32% ((D. Federman et al., 2024)). This emphasizes the clinical relevance of recognizing and treating these conditions in tandem, as they share common pathophysiological pathways.

In conclusion, the literature collectively highlights the urgent need for integrated approaches in the assessment and management of depression and MCI. Recognizing the interplay between these conditions can lead to improved outcomes through early detection, targeted interventions, and a holistic approach to patient care.

REFERENCES

1. Lara Soares Blum Malak, A., Felipe Vasconcellos, L., Santos Pereira, J., Vieira Greca, D., Cruz, M., Veiga Dias Alves, H., Sptiz, M., & Charchat-Fichman, H., 2017. Symptoms of depression in patients with mild cognitive impairment in Parkinson's disease. ncbi.nlm.nih.gov
2. Said, G., 2018. Exploring fear of dementia, subjective

- cognitive complaints and common mental health difficulties in screening for Mild Cognitive Impairment. [PDF]
3. Song, D., S.F. Yu, D., W.C. Li, P., & Sun, Q., 2019. Identifying the Factors Related to Depressive Symptoms Amongst Community-Dwelling Older Adults with Mild Cognitive Impairment. ncbi.nlm.nih.gov
 4. Han, F. F., Wang, H. X., Wu, J. J., Yao, W., Hao, C. F., & Pei, J. J., 2021. Depressive symptoms and cognitive impairment: A 10-year follow-up study from the Survey of Health, Ageing and Retirement in Europe. ncbi.nlm.nih.gov
 5. Cankaya, S., Yalciner, B., Kandemir Yilmaz, M., & Yulug, B., 2024. Differentiating MCI from depression through verbal memory scores. ncbi.nlm.nih.gov
 6. Xue, Z., Zhu, X., Wu, W., Zhu, Y., Xu, Y., & Yu, M., 2024. Synapse-Related Serum and P300 Biomarkers Predict the Occurrence of Mild Cognitive Impairment in Depression. ncbi.nlm.nih.gov
 7. D. Federman, A., Becker, J., Carnavali, F., Rivera Mindt, M., Cho, D., Pandey, G., Chan, L., Curtis, L., S. Wolf, M., & P. Wisnivesky, J., 2024. Relationship Between Cognitive Impairment and Depression Among Middle Aged and Older Adults in Primary Care. ncbi.nlm.nih.gov